

October 2, 2019

To whom it may concern:

Company Name Corporate Representative	TOHO HOLDINGS CO., LTD. Norio Hamada, Chairman of the Board and Chief Executive Officer (CEO), Representative Director (First Section of Tokyo Stock Exchange Securities Code:8129)
Contact:	Makoto Kawamura, Director and General Manager, Public and Investor Relations Department (TEL: 81-3-6838-2803)

Notice Regarding Approval of Partial Amendments to Dosage and Administration of Candesartan Tablets 2mg / 4mg / 8mg / 12mg “FFP” for KYOSOMIRAI PHARMA

TOHO HOLDINGS CO., LTD. (Headquarters: Tokyo; Chairman of the Board and Chief Executive Officer (CEO), Representative Director: Norio Hamada) is pleased to announce that KYOSOMIRAI PHARMA CO., LTD. (Headquarters: Tokyo; President and Representative Director: Atsushi Udoh), the wholly-owned subsidiary involved in pharmaceuticals manufacture and sales business, received an approval of partial amendments to dosage and administration of Candesartan Tablets 2mg / 4mg / 8mg / 12mg “FFP” on October 2, 2019 as below.

Outline of Candesartan Tablets 2mg / 4mg / 8mg / 12mg “FFP”

Class	Product Name	Original Brand Name
Long-acting angiotensin II receptor antagonist	Candesartan Tablets 2mg / 4mg / 8mg / 12mg “FFP”	BLOPRESS Tablets 2 / 4 / 8 / 12

Outline of Amendments to Dosage and Administration of Candesartan Tablets 2mg / 4mg / 8mg / 12mg “FFP”

Indications and Usage	Dosage and Administration
[Tablets 2mg/4mg/8mg/12mg] Hypertension	<u>Adults</u> The usual Candesartan dosage for adults is 4-8 mg as candesartan cilexetil once daily via oral administration and the dose can be increased up to 12 mg, if necessary. For patients with comorbid renal disorder, start with 2 mg once daily and increase up to 8 mg, if necessary. <u>Children</u> The usual Candesartan dosage for children aged 1 to under 6 years is 0.05-0.3 mg/kg as candesartan cilexetil once daily via oral administration. The usual Candesartan dosage for children aged 6 years and older is 2-8 mg as candesartan cilexetil once daily via oral administration and the dose can be increased up to 12 mg, if necessary. For patients with comorbid renal disorder, start with a low dose and increase up to 8 mg, if necessary.
[Tablets 2mg/4mg/8mg/12mg] Renal parenchymal hypertension	The usual Candesartan dosage for adults is 2 mg as candesartan cilexetil once daily via oral administration and the dose can be increased up to 8 mg, if necessary.
[Tablets 2mg/4mg/8mg] When treatment with angiotensin-converting enzyme inhibitors is unsuitable due to the condition described below: Mild to moderate chronic cardiac failure	The usual Candesartan dosage for adults is 4 mg as candesartan cilexetil once daily via oral administration and the dose can be increased up to 8 mg, if necessary. In principle, background treatment other than with an angiotensin-converting enzyme inhibitor should be continued.

(Underlined parts are revised parts.)